

On the Total Perimeter of Homothetic Convex Bodies in a Convex Container*

Adrian Dumitrescu[†] Csaba D. Tóth[‡]

August 11, 2014

Abstract

For two planar convex bodies, C and D , consider a packing S of n positive homothets of C contained in D . We estimate the total perimeter of the bodies in S , denoted $\text{per}(S)$, in terms of $\text{per}(D)$ and n . When all homothets of C touch the boundary of the container D , we show that either $\text{per}(S) = O(\log n)$ or $\text{per}(S) = O(1)$, depending on how C and D “fit together”. Apart from the constant factors, these bounds are the best possible. Specifically, we prove that $\text{per}(S) = O(1)$ if D is a convex polygon and every side of D is parallel to a corresponding segment on the boundary of C (for short, D is *parallel to C*) and $\text{per}(S) = O(\log n)$ otherwise.

When D is parallel to C but the homothets of C may lie anywhere in D , we show that $\text{per}(S) = O((1 + \text{esc}(S)) \log n / \log \log n)$, where $\text{esc}(S)$ denotes the total distance of the bodies in S from the boundary of D . Apart from the constant factor, this bound is also the best possible.

Keywords: convex body, perimeter, maximum independent set, homothet, Ford disks, traveling salesman, approximation algorithm.

1 Introduction

A finite set $S = \{C_1, \dots, C_n\}$ of convex bodies is a *packing* in a convex body (*container*) $D \subset \mathbb{R}^2$ if the bodies $C_1, \dots, C_n \in S$ are contained in D and they have pairwise disjoint interiors. The term *convex body* above refers to a compact convex set with nonempty interior in \mathbb{R}^2 . The perimeter of a convex body $C \subset \mathbb{R}^2$ is denoted $\text{per}(C)$, and the total perimeter of a packing S is denoted $\text{per}(S) = \sum_{i=1}^n \text{per}(C_i)$. Our interest is estimating $\text{per}(S)$ in terms of n . In this paper, we consider packings S that consist of positive homothets of a convex body C . A positive homothet of $C \subset \mathbb{R}^2$ is a planar set $\{\rho\mathbf{c} + \mathbf{t} : \mathbf{c} \in C\}$, where $\rho > 0$ is a scale factor and $\mathbf{t} \in \mathbb{R}^2$ is a (translation) vector. We start with an easy general bound for this case.

Proposition 1. *For every pair of convex bodies, C and D , and every packing S of n positive homothets of C in D , we have $\text{per}(S) \leq \rho(C, D)\sqrt{n}$, where $\rho(C, D)$ depends on C and D . Apart from this multiplicative constant, this bound is the best possible.*

*A preliminary version of this paper appeared in *Proceedings of the 17th International Workshop on Approximation Algorithms for Combinatorial Optimization Problems* (APPROX 2013), Berkeley, CA, 2013, LNCS 8096, pp. 96–109.

[†]Department of Computer Science, University of Wisconsin–Milwaukee, WI, USA. Email: dumitres@uwm.edu. Supported in part by NSF grant DMS-1001667.

[‡]Department of Mathematics, California State University, Northridge, Los Angeles, CA, USA; and Department of Computer Science, Tufts University, Medford, MA, USA. Email: cdtoth@acm.org.

Our goal is to derive substantially better upper bounds on $\text{per}(S)$ in terms of n in two different scenarios, motivated by applications to the traveling salesman problem with neighborhoods (TSPN). In Sections 3–4, we prove tight bounds on $\text{per}(S)$ in terms of n when all homothets in S touch the boundary of the container D (see Fig. 1). In Section 5, we prove tight bounds on $\text{per}(S)$ in terms of n and the total distance of the bodies in S from the boundary of D . Specifically, for two convex bodies, $C \subset D \subset \mathbb{R}^2$, let the *escape distance* $\text{esc}(C)$ be the distance between C and the boundary of D (Fig. 2, right); and for a packing $S = \{C_1, \dots, C_n\}$ in a container D , let $\text{esc}(S) = \sum_{i=1}^n \text{esc}(C_i)$.




Figure 1: Left: a packing of disks in a rectangle container, where all disks touch the boundary of the container. Right: a convex body C in the interior of a trapezoid D at distance $\text{esc}(C)$ from the boundary of D . The trapezoid D is *parallel to* C : every side of D is parallel and “corresponds” to a side of C .

Homothets touching the boundary of a convex container. We would like to bound $\text{per}(S)$ from above in terms of $\text{per}(D)$ and n when all homothets in S touch the boundary of D (see Fig. 1, left). Specifically, for a pair of convex bodies, C and D , let $f_{C,D}(n)$ denote the maximum perimeter $\text{per}(S)$ of a packing of n positive homothets of C in the container D , where each element of S touches the boundary of D . We would like to estimate the growth rate of $f_{C,D}(n)$ as n goes to infinity. We prove a logarithmic¹ upper bound $f_{C,D}(n) = O(\log n)$ for every pair of convex bodies, C and D .

Proposition 2. *For every pair of convex bodies, C and D , and every packing S of n positive homothets of C in D , where each element of S touches the boundary of D , we have $\text{per}(S) \leq \rho(C, D) \log n$, where $\rho(C, D)$ depends on C and D .*

The upper bound $f_{C,D}(n) = O(\log n)$ is asymptotically tight for some pairs C and D , and not so tight for others. For example, it is not hard to attain an $\Omega(\log n)$ lower bound when C is an axis-aligned square, and D is a triangle (Fig. 2, left). However, $f_{C,D}(n) = \Theta(1)$ when both C and D are axis-aligned squares. We determine $f_{C,D}(n)$ up to constant factors for all pairs of convex bodies of bounded description complexity². We start by establishing a logarithmic lower bound in the simple setting where C is a circular disk and D is a unit square.

Theorem 1. *For every $n \in \mathbb{N}$, there exists a set S of n pairwise disjoint disks lying in the unit square $U = [0, 1]^2$ and touching the boundary of U such that $\text{per}(S) = \Omega(\log n)$.*

We show that either $f_{C,D} = \Theta(\log n)$ or $f_{C,D}(n) = \Theta(1)$ depending on how C and D “fit together”. To distinguish these cases, we need the following definitions.

¹Throughout this paper, $\log x$ denotes the logarithm of x to base 2.

²A planar set has *bounded description complexity* if its boundary consists of a finite number of algebraic curves of bounded degrees.




Figure 2: Left: a square packing in a triangle where every square touches the boundary of the triangle. Right: a packing of homothetic hexagons H in a square U , where U is parallel to H and every hexagon touches the boundary of U .

Definition of “parallel” convex bodies. Denote by \mathbb{S} the set of unit vectors in \mathbb{R}^2 , that is, $\mathbb{S} = \{\mathbf{d} \in \mathbb{R}^2 : |\mathbf{d}| = 1\}$. For a vector $\mathbf{d} \in \mathbb{S}$ and a convex body C , the *supporting line* $\ell_{\mathbf{d}}(C)$ is a directed line of direction \mathbf{d} such that $\ell_{\mathbf{d}}(C)$ is tangent to C , and the closed halfplane on the left of $\ell_{\mathbf{d}}(C)$ contains C . If $\ell_{\mathbf{d}}(C) \cap C$ is a nondegenerate line segment, we refer to it as a *side* of C .

We say that a convex polygon (container) D is *parallel to* a convex body C when for every direction $\mathbf{d} \in \mathbb{S}$ if $\ell_{\mathbf{d}}(D) \cap D$ is a side of D , then $\ell_{\mathbf{d}}(C) \cap C$ is also a side of C . Figure 2 (right) depicts a square D parallel to a convex hexagon C . Note that this binary relation on convex bodies is not symmetric: it is possible that D is parallel to C , but C is not parallel to D .

Classification. We generalize the lower bound construction in Theorem 1 to arbitrary convex bodies, C and D , of bounded description complexity, where D is not parallel to C .

Theorem 2. *Let C and D be two convex bodies of bounded description complexity such that D is not parallel to C . For every $n \in \mathbb{N}$, there exists a set S of n positive homothets of C in D such that each element of S touches the boundary of D , and $\text{per}(S) \geq \rho(C, D) \log n$, where $\rho(C, D)$ depends on C and D .*

If D is a convex polygon parallel to C , and every homothet of C in a packing S of n homothets touches the boundary of D , then it is not difficult to see that $\text{per}(S)$ is bounded from above by an expression independent of n .

Proposition 3. *Let C and D be convex bodies such that D is a convex polygon parallel to C . Then every packing S of n positive homothets of C in D , where each element of S touches the boundary of D , we have $\text{per}(S) \leq \rho(C, D)$, where $\rho(C, D)$ depends on C and D .*

Total distance from the boundary of a convex container. In the general case, when the homothets of C can be in the interior of the container D , we improve the dependence on n of the general bound in Proposition 1 by using the escape distance, namely the total distance of the homothets of C from the boundary of D . The combination of Propositions 1 and 2 yields the following bound.

Proposition 4. *For every pair of convex bodies, C and D , and every packing S of n positive homothets of C in D , we have $\text{per}(S) \leq \rho(C, D)(\text{esc}(S) + \log n)$, where $\rho(C, D)$ depends on C and D .*

By Theorem 2, the logarithmic upper bound in terms of n is the best possible when D is not parallel to C . When D is a convex polygon parallel to C , we derive the following upper bound for $\text{per}(S)$, which is also asymptotically tight in terms of n .

Theorem 3. *Let C and D be two convex bodies such that D is a convex polygon parallel to C . For every packing S of n positive homothets of C in D , we have*

$$\text{per}(S) \leq \rho(C, D) (\text{per}(D) + \text{esc}(S)) \frac{\log n}{\log \log n},$$

where $\rho(C, D)$ depends on C and D . For every $n \geq 1$, there exists a packing S of n positive homothets of C in D such that

$$\text{per}(S) \geq \sigma(C, D) (\text{per}(D) + \text{esc}(S)) \frac{\log n}{\log \log n},$$

where $\sigma(C, D)$ depends on C and D .

Related previous work. We consider the total perimeter $\text{per}(S)$ of a packing S of n homothets of a convex body C in a convex container D in Euclidean plane. Other variants have also been considered: (1) If S is a packing of n arbitrary convex bodies in D , then it is easy to subdivide D by $n - 1$ near diameter segments into n convex bodies of total perimeter close to $\text{per}(D) + 2(n - 2)\text{diam}(D)$. Glazyrin and Morić [11] have recently proved that this lower bound is the best possible when D is a square or a triangle. For an arbitrary convex body D , they prove an upper bound of $\text{per}(S) \leq 1.22 \text{per}(D) + 2(n - 1)\text{diam}(D)$. (2) If all bodies in S are congruent to a convex body C , then $\text{per}(S) = n \text{per}(C)$, and bounding $\text{per}(S)$ from above reduces to the classic problem of determining the maximum number of interior-disjoint congruent copies of C that fit in D [5, Section 1.6].

Considerations of the total surface area of a ball packing in \mathbb{R}^3 also play an important role in a strong version of the Kepler conjecture [3, 13].

Motivation. In the *Euclidean Traveling Salesman Problem* (ETSP), given a set S of n points in \mathbb{R}^d , one wants to find a closed polygonal chain (*tour*) of minimum Euclidean length whose vertex set is S . The Euclidean TSP is known to be NP-hard, but it admits a PTAS in \mathbb{R}^d , where $d \in \mathbb{N}$ is constant [2]. In the *TSP with Neighborhoods* (TSPN), given a set of n sets (neighborhoods) in \mathbb{R}^d , one wants to find a closed polygonal chain of minimum Euclidean length that has a vertex in each neighborhood. The neighborhoods are typically simple geometric objects (of bounded description complexity) such as disks, rectangles, line segments, or lines. While TSPN is known to be NP-hard, it admits a PTAS for certain types of neighborhoods [16], but is hard to approximate for others [6].

For n connected (possibly overlapping) neighborhoods in the plane, TSPN can be approximated with ratio $O(\log n)$ by an algorithm of Mata and Mitchell [15]. See also the survey by Bern and Eppstein [4] for a short outline of this algorithm. At its core, the $O(\log n)$ -approximation relies on the following early result by Levkopoulos and Lingas [14]: every (simple) rectilinear polygon P with n vertices, r of which are reflex, can be partitioned into rectangles of total perimeter $O(\text{per}(P) \log r)$ in $O(n \log n)$ time.

A natural approach for finding a solution to TSPN is the following [7, 9] (in particular, it achieves a constant-ratio approximation for unit disks): Given a set S of n neighborhoods, compute a maximal subset $I \subseteq S$ of pairwise disjoint neighborhoods (i.e., a packing), compute a good tour for I , and then augment it by traversing the boundary of each set in I . Since each neighborhood in $S \setminus I$ intersects some neighborhood in I , the augmented tour visits all members of S . This approach is particularly appealing since good approximation algorithms are often available for pairwise disjoint neighborhoods [16]. The bottleneck of this approach is the length increase incurred by extending a tour of I by the total perimeter of the neighborhoods in I . An upper bound

$\text{per}(I) = o(\text{OPT}(I) \log n)$ would immediately imply an improved $o(\log n)$ -factor approximation ratio for TSPN.

Theorem 2 shows that this approach cannot beat the $O(\log n)$ approximation ratio for most types of neighborhoods (e.g., circular disks). In the current formulation, Proposition 2 yields the upper bound $\text{per}(I) = O(\log n)$ assuming a convex container, so in order to use this bound, a tour of I needs to be augmented into a convex partition; this may increase the length by a $\Theta(\log n / \log \log n)$ -factor in the worst case [8, 14]. For convex polygonal neighborhoods, the bound $\text{per}(I) = O(1)$ in Proposition 3 is applicable after a tour for I has been augmented into a convex partition with *parallel* edges (e.g., this is possible for axis-aligned rectangle neighborhoods, and an axis-aligned approximation of the optimal tour for I). The convex partition of a polygon with $O(1)$ distinct orientations, however, may increase the length by a $\Theta(\log n)$ -factor in the worst case [14]. Overall our results show that we cannot beat the current $O(\log n)$ ratio for TSPN for any type of homothetic neighborhoods if we start with an arbitrary independent set I and an arbitrary near-optimal tour for I .

2 Preliminaries: A Few Easy Pieces

Proof of Proposition 1. Let $\mu_i > 0$ denote the homothety factor of C_i , i.e., $C_i = \mu_i C$, for $i = 1, \dots, n$. Since S is a packing we have $\sum_{i=1}^n \mu_i^2 \text{area}(C) \leq \text{area}(D)$. By the Cauchy-Schwarz inequality we have $(\sum_{i=1}^n \mu_i)^2 \leq n \sum_{i=1}^n \mu_i^2$. It follows that

$$\begin{aligned} \text{per}(S) &= \sum_{i=1}^n \text{per}(C_i) = \text{per}(C) \sum_{i=1}^n \mu_i \\ &\leq \text{per}(C) \sqrt{n} \sqrt{\left(\sum_{i=1}^n \mu_i^2 \right)} \leq \text{per}(C) \sqrt{\frac{\text{area}(D)}{\text{area}(C)}} \sqrt{n}. \end{aligned}$$

Set now $\rho(C, D) := \text{per}(C) \sqrt{\text{area}(D) / \text{area}(C)}$, and the proof of the upper bound is complete.

For the lower bound, consider two convex bodies, C and D . Let U be a maximal axis-aligned square inscribed in D , and let μC be the largest positive homothet of C that fits into U . Note that $\mu = \mu(C, D)$ is a constant that depends on C and D only. Subdivide U into $\lceil \sqrt{n} \rceil^2$ congruent copies of the square $\frac{1}{\lceil \sqrt{n} \rceil} U$. Let S be the packing of n translates of $\frac{\mu}{\lceil \sqrt{n} \rceil} C$, with at most one in each square $\frac{1}{\lceil \sqrt{n} \rceil} U$. The total perimeter of the packing is $\text{per}(S) = n \cdot \frac{\mu}{\lceil \sqrt{n} \rceil} \text{per}(C) = \Theta(\sqrt{n})$, as claimed. \square

Proof of Proposition 2. Let $S = \{C_1, \dots, C_n\}$ be a packing of n homothets of C in D where each element of S touches the boundary of D . Observe that $\text{per}(C_i) \leq \text{per}(D)$ for all $i = 1, \dots, n$. Partition the elements of S into subsets as follows. For $k = 1, \dots, \lceil \log n \rceil$, let S_k denote the set of homothets C_i such that $\text{per}(D)/2^k < \text{per}(C_i) \leq \text{per}(D)/2^{k-1}$; and let S_0 be the set of homothets C_i of perimeter less than $\text{per}(D)/2^{\lceil \log n \rceil}$. Then the sum of perimeters of the elements in S_0 is $\text{per}(S_0) \leq n \text{per}(D)/2^{\lceil \log n \rceil} \leq \text{per}(D)$, since $S_0 \subseteq S$ contains at most n elements altogether.

For $k = 1, \dots, \lceil \log n \rceil$, the diameter of each $C_i \in S_k$ is bounded above by

$$\text{diam}(C_i) < \text{per}(C_i)/2 \leq \text{per}(D)/2^k. \quad (1)$$

Consequently, every point of a body $C_i \in S_k$ lies at distance at most $\text{per}(D)/2^k$ from the boundary of D , denoted ∂D . Let R_k be the set of points in D at distance at most $\text{per}(D)/2^k$ from ∂D . Then

$$\text{area}(R_k) \leq \text{per}(D) \frac{\text{per}(D)}{2^k} = \frac{(\text{per}(D))^2}{2^k}. \quad (2)$$

Since S consists of homothets, the area of any element $C_i \in S_k$ is bounded from below by

$$\text{area}(C_i) = \text{area}(C) \left(\frac{\text{per}(C_i)}{\text{per}(C)} \right)^2 \geq \text{area}(C) \left(\frac{\text{per}(D)}{2^k \text{per}(C)} \right)^2. \quad (3)$$

By a volume argument, (2) and (3) yield

$$|S_k| \leq \frac{\text{area}(R_k)}{\min_{C_i \in S_k} \text{area}(C_i)} \leq \frac{(\text{per}(D))^2/2^k}{\text{area}(C)(\text{per}(D))^2/(2^k \text{per}(C))^2} = \frac{(\text{per}(C))^2}{\text{area}(C)} 2^k.$$

Since for $C_i \in S_k$, $k = 1, \dots, \lceil \log n \rceil$, we have $\text{per}(C_i) \leq \text{per}(D)/2^{k-1}$, it follows that

$$\text{per}(S_k) \leq |S_k| \cdot \frac{\text{per}(D)}{2^{k-1}} \leq 2 \frac{(\text{per}(C))^2}{\text{area}(C)} \text{per}(D).$$

Hence the sum of perimeters of all elements in S is bounded by

$$\text{per}(S) = \sum_{k=0}^{\lceil \log n \rceil} \text{per}(S_k) \leq \left(1 + 2 \frac{(\text{per}(C))^2}{\text{area}(C)} \lceil \log n \rceil \right) \text{per}(D),$$

as required. \square

Proof of Proposition 3. Let $\rho'(C)$ denote the ratio between $\text{per}(C)$ and the length of a shortest side of C . Recall that each $C_i \in S$ touches the boundary of polygon D . Since D is parallel to C , the side of D that supports C_i must contain a side of C_i . Let a_i denote the length of this side.

$$\text{per}(S) = \sum_{i=1}^n \text{per}(C_i) = \sum_{i=1}^n a_i \frac{\text{per}(C_i)}{a_i} \leq \rho'(C) \sum_{i=1}^n a_i \leq \rho'(C) \text{per}(D).$$

Set now $\rho(C, D) := \rho'(C) \text{per}(D)$, and the proof is complete. \square

Proof of Proposition 4. The proof is similar to that of Proposition 2 with a few adjustments. Let $S = \{C_1, \dots, C_n\}$ be a packing of n homothets of C in D . Note that $\text{per}(C_i) \leq \text{per}(D)$ for all $i = 1, \dots, n$. Partition the elements of S into subsets as follows. Let

$$S^{\text{in}} = \{C_i \in S : \text{per}(C_i) \leq \text{esc}(C_i)\} \text{ and } S^{\text{bd}} = S \setminus S^{\text{in}}.$$

For $k = 1, \dots, \lceil \log n \rceil$, let S_k denote the set of homothets $C_i \in S^{\text{bd}}$ such that $\text{per}(D)/2^k < \text{per}(C_i) \leq \text{per}(D)/2^{k-1}$; and let S_0 be the set of homothets $C_i \in S^{\text{bd}}$ of perimeter at most $\text{per}(D)/2^{\lceil \log n \rceil}$.

The sum of perimeters of the elements in S^{in} is $\text{per}(S^{\text{in}}) \leq \text{esc}(S^{\text{in}}) \leq \text{esc}(S)$. We next consider the elements in S^{bd} . The sum of perimeters of the elements in S_0 is $\text{per}(S_0) \leq n \text{per}(D)/2^{\lceil \log n \rceil} \leq \text{per}(D)$, since $S_0 \subseteq S$ contains at most n elements altogether.

For $k = 1, \dots, \lceil \log n \rceil$, the diameter of each $C_i \in S_k$ is bounded above by $\text{diam}(C_i) < \text{per}(C_i)/2 \leq \text{per}(D)/2^k$. Observe that every point of a body $C_i \in S_k$ lies at distance at most $\text{esc}(C_i) + \text{diam}(C_i) \leq \text{per}(C_i) + \text{diam}(C_i) \leq 1.5 \text{per}(C_i) \leq 3 \text{per}(D)/2^k$ from ∂D . Let now R_k be the set of points in D at distance at most $3 \text{per}(D)/2^k$ from ∂D . Then

$$\text{area}(R_k) \leq \text{per}(D) \frac{3 \text{per}(D)}{2^k} = \frac{3 (\text{per}(D))^2}{2^k}.$$

Analogously to the proof of Proposition 2, a volume argument yields

$$|S_k| \leq 3 \frac{(\text{per}(C))^2}{\text{area}(C)} 2^k.$$

It follows that

$$\text{per}(S_k) \leq |S_k| \cdot \frac{\text{per}(D)}{2^{k-1}} \leq 6 \frac{(\text{per}(C))^2}{\text{area}(C)} \text{per}(D).$$

Hence the sum of perimeters of all elements in S is bounded by

$$\text{per}(S) \leq \text{esc}(S) + \left(1 + 6 \frac{(\text{per}(C))^2}{\text{area}(C)} \lceil \log n \rceil\right) \text{per}(D),$$

as required. □

3 Disks Touching the Boundary of a Square: Proof of Theorem 1

We show that there exists a packing of $O(n)$ disks in the unit square U such that every disk touches the x -axis, and the sum of their diameters is $\Omega(\log n)$. We present two constructions attaining this bound: (i) an *explicit* construction in Subsection 3.1 which will be generalized in Section 4; and (ii) a *greedy* disk packing.

3.1 An Explicit Construction

For convenience, we use the unit square $[-\frac{1}{2}, \frac{1}{2}] \times [0, 1]$ for our construction. To each disk we associate its vertical *projection interval* (on the x -axis). The algorithm greedily chooses disks of monotonically decreasing radii such that (1) every diameter is $1/8^k$ for some $k \in \mathbb{N}$; and (2) if the projection intervals of two disks overlap, then one interval contains the other.

For $k = 0, 1, \dots, \lceil \log_8 n \rceil$, denote by S_k the set of disks of diameter $1/8^k$, constructed by our algorithm. We recursively allocate a finite union of intervals $X_k \subset [-\frac{1}{2}, \frac{1}{2}]$ to S_k , and then choose disks in S_k such that their projection intervals lie in X_k . Initially, $X_0 = [-\frac{1}{2}, \frac{1}{2}]$, and S_0 contains the disk of diameter 1 inscribed in $[-\frac{1}{2}, \frac{1}{2}] \times [0, 1]$. The length of each maximal interval $I \subseteq X_k$ will be a multiple of $1/8^k$, so I can be covered by projection intervals of interior-disjoint disks of diameter $1/8^k$ touching the x -axis. Every interval $I \subseteq X_k$ will have the property that any disk of diameter $1/8^k$ whose projection interval is in I is disjoint from any (larger) disk in S_j , $j < k$.

Consider the disk Q of diameter 1, centered at $(0, \frac{1}{2})$, and tangent to the x -axis (see Fig. 3). It can be easily verified that:

- (i) the locus of centers of disks tangent to both Q and the x -axis is the parabola $y = \frac{1}{2}x^2$; and
- (ii) any disk of diameter $1/8$ and tangent to the x -axis whose projection interval is in $I_1(Q) = [-\frac{1}{2}, -\frac{1}{4}] \cup [\frac{1}{4}, \frac{1}{2}]$ is disjoint from Q .

Indeed, the center of any such disk is $(x_1, \frac{1}{16})$, for $x_1 \leq -5/16$ or $x_1 \geq 5/16$, and hence lies below the parabola $y = \frac{1}{2}x^2$. Similarly, for all $k \in \mathbb{N}$, any disk of diameter $1/8^k$ and tangent to the x -axis whose projection interval is in $I_k(Q) = [-\frac{1}{2^k}, -\frac{1}{2^{k+1}}] \cup [\frac{1}{2^{k+1}}, \frac{1}{2^k}]$ is disjoint from Q . We can extend this definition to an arbitrary disk D tangent to the x -axis, by appropriate scaling and translation of the intervals. If D has radius r and projection interval $[x - \frac{w}{2}, x + \frac{w}{2}]$, then let $I_0(D) = [x - \frac{w}{2}, x + \frac{w}{2}]$, and for an integer $k \geq 1$, denote by $I_k(D) = [x - \frac{w}{2^k}, x - \frac{w}{2^{k+1}}] \cup [x + \frac{w}{2^{k+1}}, x + \frac{w}{2^k}]$ the pair of intervals corresponding to $I_k(Q)$.




Figure 3: Disk Q and the exponentially decreasing pairs of intervals $I_k(Q)$, $k = 1, 2, \dots$

We can now recursively allocate intervals in X_k and choose disks in S_k ($k = 0, 1, \dots, \lfloor \log_8 n \rfloor$) as follows. Recall that $X_0 = [-\frac{1}{2}, \frac{1}{2}]$, and S_0 contains a single disk of unit diameter inscribed in the unit square $[-\frac{1}{2}, \frac{1}{2}] \times [0, 1]$. Assume that we have already defined the union of intervals X_{k-1} , and selected disks in S_{k-1} . Let X_k be the union of the interval pairs $I_{k-j}(D)$ for all $D \in S_j$ and $j = 0, 1, \dots, k-1$. Place the maximum number of disks of diameter $1/8^k$ into S_k such that their projection intervals are contained in X_k . For a disk $D \in S_j$ ($j = 0, 1, \dots, k-1$) of diameter $1/8^j$, the two intervals in X_{k-j} each have length $\frac{1}{2} \cdot \frac{1}{2^{k-j}} \cdot \frac{1}{8^j} = \frac{8^{k-j}}{2} \cdot \frac{1}{8^k}$, so they can each accommodate the projection intervals of $\frac{8^{k-j}}{2}$ disks in S_k .

We prove by induction on k that the length of X_k is $\frac{1}{2}$, and so the sum of the diameters of the disks in S_k is $\frac{1}{2}$, for $k = 1, 2, \dots, \lfloor \log_8 n \rfloor$. The interval $X_0 = [-\frac{1}{2}, \frac{1}{2}]$ has length 1. The union of intervals $X_1 = [-\frac{1}{2}, -\frac{1}{4}] \cup [\frac{1}{4}, \frac{1}{2}]$ has length $\frac{1}{2}$. For $k = 2, \dots, \lfloor \log_8 n \rfloor$, the union X_k consists of two types of (disjoint) intervals: (a) The pair of intervals $I_1(D)$ for every $D \in S_{k-1}$ covers half of the projection interval of D . Over all $D \in S_{k-1}$, they jointly cover half the length of X_{k-1} . (b) Each pair of intervals $I_{k-j}(D)$ for $D \in S_j$, $j = 0, \dots, k-2$, has half the length of $I_{k-j-1}(D)$. So the sum of the lengths of these intervals is half the length of X_{k-1} , although these intervals are disjoint from X_{k-1} . Altogether, the sum of lengths of all intervals in X_k is the same as the length of X_{k-1} . By induction, the length of X_{k-1} is $\frac{1}{2}$, hence the length of X_k is also $\frac{1}{2}$, as claimed. This immediately implies that the sum of diameters of the disks in $\bigcup_{k=0}^{\lfloor \log_8 n \rfloor} S_k$ is $1 + \frac{1}{2} \lfloor \log_8 n \rfloor$. Finally, one can verify that the total number of disks used is $O(n)$. Write $K = \lfloor \log_8 n \rfloor$. Indeed, $|S_0| = 1$, and $|S_k| = |X_k|/8^{-k} = 8^k/2$, for $k = 1, \dots, K$, where $|X_k|$ denotes the total length of the intervals in X_k . Consequently, $|S_0| + \sum_{k=1}^K |S_k| = O(8^K) = O(n)$, as required. \square

3.2 A Greedy Disk Packing

The following simple greedy algorithm produces a packing S_n of n disks in the unit square $U = [0, 1]^2$ with all disks touching the boundary of U and whose total perimeter is $\Omega(\log n)$. For $i = 1$ to n , let C_i be a disk of maximum radius that lies in $U \setminus (\bigcup_{j < i} C_j)$ and intersects ∂U , and let $S_n = \{C_1, \dots, C_n\}$; refer to Fig. 4 (left). The radius of C_1 is $1/2$, the radii of C_2, \dots, C_5 are $3 - 2\sqrt{2}$, etc. We use Apollonian circle packings [12] to derive the lower bound $\text{per}(S_n) = \Omega(\log n)$.

We now consider a greedy algorithm in a slightly different setting. For $r_1, r_2 > 0$, we construct a set $F_n(r_1, r_2)$ of n disks by the following greedy algorithm. Let A_1 and A_2 be two tangent disks of radii r_1 and r_2 that are also tangent to the x -axis from above. Let I be the horizontal segment

between the tangency points of A_1 and A_2 with the x -axis. For $i = 3, \dots, n$, let A_i be the disk of maximum radius tangent to segment I , lying above the x -axis, and disjoint from the interior of all disks A_j , $j < i$. See Fig. 4 (right), where $r_1 = r_2 = 1/2$. We now compare the total perimeter of the two greedy disk packings described above.




Figure 4: Left: A greedy packing of $n = 7$ disks in $[0, 1]^2$. Right: Ford disks visible in the window $[0, 1]^2$.

Proposition 5. $\text{per}(S_n) \geq \text{per}(F_n(1/2, 3 - 2\sqrt{2}))$.

Proof. Recall that the first two disks in S_n have radii $1/2$ and $3 - 2\sqrt{2}$, respectively. Let I be the line segment between the tangency points of A_1 and A_2 with the bottom side of $[0, 1]^2$. Because of the greedy strategy, all disks in S_n that touch the segment I are in $F_n(1/2, 3 - 2\sqrt{2})$. The radius of every disk in $S_n \setminus F_n(1/2, 3 - 2\sqrt{2})$ is at least as large as any disk in $F_n(1/2, 3 - 2\sqrt{2}) \setminus S_n$. Therefore, there is a one-to-one correspondence between S_n and $F_n(1/2, 3 - 2\sqrt{2})$ such that each disk in S_n corresponds to a disk of the same or smaller radius in $F_n(1/2, 3 - 2\sqrt{2})$. \square

Given two tangent disks of radii r_1 and r_2 that are also tangent to the x -axis, there is a unique disk tangent to both these disks and the x -axis, and its radius r_3 satisfies $r_3^{-1/2} = r_1^{-1/2} + r_2^{-1/2}$. Observe that $r_3 = r_3(r_1, r_2)$ is a continuous and monotonically increasing function of both variables, r_1 and r_2 . Therefore, if $r_1 \leq r'_1$ and $r_2 \leq r'_2$, then

$$\text{per}(F_n(r_1, r_2)) \leq \text{per}(F_n(r'_1, r'_2)). \quad (4)$$

This observation allows us to bound $\text{per}(S_n)$ from below by the perimeter of a finite subfamily of Ford disks [10]: this is a packing of an *infinite* set of disks in the halfplane $\{(x, y) \in \mathbb{R}^2 : y \geq 0\}$, where each disk is tangent to the x -axis from above. Every pair $(p, q) \in \mathbb{N}^2$ of relative prime positive integers, the Ford disk $C_{p,q}$ is of radius $1/(2q^2)$ centered at $(p/q, 1/(2q^2))$; see Fig. 4 (right). The Ford disks have pairwise disjoint interiors [10]. The Ford disks $C_{p,1}$ have the largest radius $1/2$; all other Ford disks have smaller radii and each is tangent to two larger Ford disks [10]. Hence, the set of the n largest Ford disks that touch the unit segment $[0, 1]$ is exactly $F_n(1/2, 1/2)$.

Proposition 6. $\text{per}(F_n(1/2, 1/2)) = \Omega(\log n)$.

Proof. For a positive integer Q , the number of Ford disks of radius at least $\frac{1}{2Q^2}$ touching the unit segment $[0, 1]$ is $1 + \sum_{q=1}^Q \varphi(q)$, where $\varphi(\cdot)$ is Euler's totient function, i.e., the number positive integers less than or equal to q that are relatively prime to q . It is known [1, Theorem 3.7, p. 62] that

$$\sum_{q=1}^Q \varphi(q) = \frac{3}{\pi^2} Q^2 + O(Q \log Q).$$

Hence, for a suitably large $Q = \Theta(\sqrt{n})$, there exists exactly n Ford disks of radius at least $\frac{1}{2Q^2}$ that touch $[0, 1]$. Let $F_n(1/2, 1/2)$ be the subset of these n Ford disks. Then we have

$$\text{per}(F_n) = \sum_{q=1}^Q \varphi(q) \cdot 2\pi \cdot \frac{1}{2q^2} = \pi \sum_{q=1}^Q \frac{\varphi(q)}{q^2}.$$

It is also known [1, Exercise 6, p. 71] that

$$\sum_{q=1}^Q \frac{\varphi(q)}{q^2} = \frac{6}{\pi^2} \ln Q + O\left(\frac{\log Q}{Q}\right). \quad (5)$$

Using this estimate, we have

$$\text{per}(F_n) = \pi \left(\sum_{q=1}^Q \frac{\varphi(q)}{q^2} \right) = \Omega(\log Q) = \Omega(\log \sqrt{n}) = \Omega(\log n),$$

as claimed. □

The bounds in Propositions 5-6 in conjunction with (4) yield

$$\begin{aligned} \text{per}(S_n) &\geq \text{per}(F_n(1/2, 3 - 2\sqrt{2})) \geq \text{per}(F_n(3 - 2\sqrt{2}, 3 - 2\sqrt{2})) \\ &= \Omega(\text{per}(F_n(1/2, 1/2))) = \Omega(\log n). \end{aligned}$$

When C is a disk and the container D is any other convex body, the above argument goes through and shows that a greedy packing S_n has total perimeter $\text{per}(S) = \Omega(\log n)$, where the constant of proportionality depends on D . However, when C is not a circular disk, the theory of Apollonian circles does not apply.

4 Homothets Touching the Boundary: Proof of Theorem 2

We construct a packing S of perimeter $\text{per}(S) = \Omega(\log n)$ for given C and D . Let C and D be two convex bodies with bounded description complexity. We wish to argue analogously to the case of disks in a square. Therefore, we choose an arc $\gamma \subset \partial D$ that is smooth and sufficiently “flat,” but contains no side parallel to a corresponding side of C . Then we build a hierarchy of homothets of C touching the arc γ , so that the depth of the hierarchy is $O(\log n)$, and the homothety factors decrease by a constant between two consecutive levels.

We choose an arc $\gamma \subset \partial D$ as follows. If D has a side with some direction $\mathbf{d} \in \mathbb{S}$ such that C has no parallel side of the same direction \mathbf{d} , then let γ be this side of D . Otherwise, ∂D contains an algebraic curve γ_1 of degree 2 or higher. Let $q \in \gamma_1$ be an interior point of this curve such that γ_1 is twice differentiable at q . Assume, after a rigid transformation of D if necessary, that $q = (0, 0)$ is the origin and the supporting line of D at q is the x -axis. By the inverse function theorem, there is an arc $\gamma_2 \subseteq \gamma_1$, containing q , such that γ_2 is the graph of a twice differentiable function of x . Finally, let $\gamma \subset \gamma_2$ be an arc such that the part of ∂C that has the same tangent lines as γ contains no segments (sides).

For every point $p \in \gamma$, let $p = (x_p, y_p)$, and let s_p be the slope of the tangent line of D at p . Then the tangent line of D at $p \in \gamma$ is $\ell_p(x) = s_p(x - x_p)$. For any positive homothet ρC of C , let $(\rho C)_p$ denote a translate of ρC tangent to ℓ_p at point p (Fig. 5). If both C and D have bounded




Figure 5: If a homothet C_p is tangent to $\gamma \subset \partial D$ at point p , then there are polynomials α_p and β_p that separate γ from C_p . We can place a constant number of congruent homothets of C between α_p and β_p whose vertical projections cover $I_1(Q)$. These homothets can be translated vertically down to touch γ .

description complexity, then there are constants $\rho_0 > 0$, $\kappa \in \mathbb{N}$ and $A < B$, such that for every point $p \in \gamma$ and every homothety factor ρ , $0 < \rho < \rho_0$, the polynomials

$$\alpha_p(x) = A|x - x_p|^\kappa + s_p(x - x_p), \quad \text{and} \quad \beta_p(x) = B|x - x_p|^\kappa + s_p(x - x_p),$$

separate γ from the convex body $(\rho C)_p$, that is, for every vertical line $\ell : x = x_0$, any intersection point $\ell \cap \gamma$ is at or below $(x_0, \alpha(x_0))$, and any intersection point in $\ell \cap (\rho C)_p$ is at or above $(x_0, \beta(x_0))$.

Similarly to the proof of Theorem 1, the construction is guided by nested *projection intervals*. Let $p \in \gamma$ be the midpoint of γ , and let $Q = (\rho C)_p$ for a sufficiently small $0 < \rho < \rho_0$ such that $Q \subseteq D$ and the vertical projection interval $I(Q)$ of Q is contained in the vertical projection of γ . Note that $Q = (\rho C)_p$ is tangent to γ at point $p \in \gamma$, since $0 < \rho < \rho_0$. For $k = 0, 1, \dots$, we recursively define disjoint intervals or interval pairs $I_k(Q) \subset I(Q)$ of length $|I_k(Q)| = |I(Q)|/2^k$, starting with $I_0(Q) = I(Q)$. During the recursion, we maintain the invariant that the set $J_k(Q) = I(Q) \setminus \bigcup_{j < k} I_j(Q)$ is an interval of length $|I(Q)|/2^{k-1}$ that contains x_p . Assume that $I_0(Q), \dots, I_{k-1}(Q)$ have been defined, and we need to choose $I_k(Q) \subset J_k(Q)$. Divide the interval $J_k(Q)$ into three closed intervals: a middle interval of length $\frac{1}{4}|J_k(Q)|$, and a left and a right interval, each of length $\frac{3}{8}|J_k(Q)|$. If x_p lies in the middle interval of $J_k(Q)$, then let $I_k(Q)$ be a pair of intervals that consists of the left and right *quarters* of $J_k(Q)$. If x_p lies in the left (resp., right) third $J_k(Q)$, then let $I_k(Q)$ be the right (resp., left) *half* of $J_k(Q)$.

It is now an easy matter to check (by induction on k) that $|x - x_p| \geq |I(Q)|/8^k$ for all $x \in I_k$. Consequently,

$$\beta_p(x) - \alpha_p(x) \geq (B - A) \cdot \left(\frac{|I(Q)|}{8^k} \right)^\kappa \quad (6)$$

for all $x \in I_k(Q)$. There is a constant $\mu > 0$ such that a homothet $\mu^k Q$ with arbitrary projection interval in $I_k(Q)$ fits between the curves α_p and β_p . Refer to Fig. 5. Therefore we can populate the region between the curves α_p and β_p and above $I_k(Q)$ with homothets of $\rho'(Q)$, of homothety factors $\mu^k/2 < \rho' \leq \mu^k$, such that their projection intervals are pairwise disjoint and cover $I_k(Q)$. By translating these homothets vertically until they touch γ , they remain disjoint from Q and preserve their projection intervals. We can now repeat the construction of the previous section and obtain $\lceil \log_{(2/\mu)} n \rceil$ layers of homothets touching γ , such that the total length of the projections of the homothets in each layer is $\Theta(1)$. Consequently, the total perimeter of the homothets in each layer is $\Theta(1)$, and the overall perimeter of the packing is $\Omega(\log n)$, as required. \square

5 Bounds in Term of the Escape Distance: Proof of Theorem 3

Upper bound. Let $S = \{C_1, \dots, C_n\}$ be a packing of n homothets of a convex body C in a container D such that D is a convex polygon parallel to C . For each element $C_i \in S$, $\text{esc}(C_i)$ is the distance between a side of D and a corresponding side of C_i . For each side a of D , let $S_a \subseteq S$ denote the set of $C_i \in S$ for which a is the closest side of D (ties are broken arbitrarily). Since D has finitely many sides, it is enough to show that for each side a of D , we have

$$\text{per}(S_a) \leq \rho_a(C, D) (\text{per}(D) + \text{esc}(S)) \frac{\log |S_a|}{\log \log |S_a|},$$

where $\rho_a(C, D)$ depends on a , C and D only.

Suppose that $S_a = \{C_1, \dots, C_n\}$ is a packing of n homothets of C such that $\text{esc}(C_i)$ equals the distance between C_i and side a of D . Assume for convenience that a is horizontal. Let $c \subset \partial C$ be the side of C corresponding to the side a of D . Let $\rho_1 = \text{per}(C)/|c|$, and then we can write $\text{per}(C) = \rho_1|c|$.

Denote by $b \subset c$ the line segment of length $|b| = |c|/2$ with the same midpoint as c . Refer to Fig. 6 (left). Since C is a convex body, the two vertical lines through the two endpoints of b intersect C in two line segments denoted h_1 and h_2 , respectively. Let $\rho_2 = \min(|h_1|, |h_2|)/|b|$, and then $\min(|h_1|, |h_2|) = \rho_2|b|$. By convexity, every vertical line that intersects segment b intersects C in a vertical segment of length at least $\rho_2|b|$. Note that ρ_1 and ρ_2 are constants depending on C and D . For each homothet $C_i \in S_a$, let $b_i \subset \partial C_i$ be the homothetic copy of segment $b \subset \partial C$.




Figure 6: Left: A convex body C with a horizontal side c . The segment $b \subset c$ has length $|b| = |c|/2$, and the vertical segments h_1 and h_2 are incident to the endpoints of b . Right: Two homothets, C_i and C_j , in a convex container D . The vertical projections of b_i and b_j onto the horizontal side a are proj_i and proj_j .

Put $\lambda = 2\lceil \log n / \log \log n \rceil$. Partition S_a into two subsets $S_a = S_{\text{far}} \cup S_{\text{close}}$ as follows. For each $C_i \in S_a$, let $C_i \in S_{\text{close}}$ if $\text{esc}(C_i) < \rho_2|b_i|/\lambda$, and $C_i \in S_{\text{far}}$ otherwise. For each homothet $C_i \in S_{\text{close}}$, let $\text{proj}_i \subseteq a$ denote the vertical projection of segment b_i onto the horizontal side a (refer to Fig. 6, right). The perimeter of each $C_i \in S_a$ is $\text{per}(C_i) = \rho_1|c_i| = 2\rho_1|b_i| = 2\rho_1|\text{proj}_i|$. We have

$$\text{per}(S_{\text{far}}) = \sum_{C_i \in S_{\text{far}}} \text{per}(C_i) = \sum_{C_i \in S_{\text{far}}} 2\rho_1|b_i| \leq \sum_{C_i \in S_{\text{far}}} 2\rho_1 \frac{\text{esc}(C_i) \lambda}{\rho_2} \leq \frac{2\rho_1 \text{esc}(S)}{\rho_2} \lambda. \quad (7)$$

It remains the estimate $\text{per}(S_{\text{close}})$ as an expression of λ .

$$\sum_{C_i \in S_{\text{close}}} \text{per}(C_i) = 2\rho_1 \sum_{C_i \in S_{\text{close}}} |\text{proj}_i|. \quad (8)$$

Define the *depth* function for every point of the horizontal side a by

$$d : a \rightarrow \mathbb{N}, \quad d(x) = |\{C_i \in S_{\text{close}} : x \in \text{proj}_i\}|.$$

That is, $d(x)$ is the number of homothets such that the vertical projection of segment b_i contains point x . For every positive integer $k \in \mathbb{N}$, let

$$I_k = \{x \in a : d(x) \geq k\},$$

that is, I_k is the set of points of depth at least k . Since S_{close} is finite, the set $I_k \subseteq a$ is measurable. Denote by $|I_k|$ the measure (total length) of I_k . By definition, we have $|a| \geq |I_1| \geq |I_2| \geq \dots$. A standard double counting for the integral $\int_{x \in a} d(x) dx$ yields

$$\sum_{C_i \in S_{\text{close}}} |\text{proj}_i| = \sum_{k=1}^{\infty} |I_k|. \quad (9)$$

If $d(x) = k$ for some point $x \in a$, then k segments b_i lie above x . Each $C_i \in S_{\text{close}}$ is at distance $\text{esc}(C_i) < \rho_2 |b_i|/\lambda$ from a . Suppose that proj_i and proj_j intersect for $C_i, C_j \in S_{\text{close}}$ (Fig. 6, right). Then one of them has to be closer to a than the other: we may assume w.l.o.g. $\text{esc}(C_j) < \text{esc}(C_i)$. Now a vertical segment between $b_i \subset C_i$ and $\text{proj}_i \subset a$ intersects b_j . The length of this segment, $\text{esc}(C_i)$, satisfies $\rho_2 |b_j| \leq \text{esc}(C_i) < \rho_2 |b_i|/\lambda$. Consequently, $|b_j| < |b_i|/\lambda$ (or, equivalently, $|\text{proj}_j| < |\text{proj}_i|/\lambda$) holds for any consecutive homothets above point $x \in a$. In particular, if proj_i is the k th largest projection containing $x \in a$, then $|\text{proj}_i| \leq |a|/\lambda^{k-1} = |a|\lambda^{1-k}$.

We claim that

$$|I_k| \leq |a|\lambda^{\lambda-k} \quad \text{for } k \geq \lambda + 1. \quad (10)$$

Suppose, to the contrary, that $|I_k| > |a|\lambda^{\lambda-k}$ for some $k \geq \lambda + 1$. Then there are homothets $C_i \in S_{\text{close}}$ with $|\text{proj}_i| \leq |a|/\lambda^{k-1}$ that jointly project to I_k . Assuming that $|I_k| > |a|\lambda^{\lambda-k}$, it follows that the number of these homothets is at least

$$\frac{|a|\lambda^{\lambda-k}}{|a|\lambda^{1-k}} = \lambda^{\lambda-1} = \left(2 \left\lceil \frac{\log n}{\log \log n} \right\rceil\right)^{2^{\lceil \frac{\log n}{\log \log n} \rceil - 1}} > n, \quad (11)$$

for $n \geq 3$ (recall that $x^x = n$ solves to $x = \Theta(\log n / \log \log n)$). This contradicts the fact that $S_{\text{close}} \subseteq S$ has at most n elements. Combining (8), (9), and (10), we conclude that

$$\begin{aligned} \text{per}(S_{\text{close}}) &= 2\rho_1 \sum_{k=1}^{\infty} |I_k| \leq 2\rho_1 \left(\lambda |I_1| + \sum_{k=\lambda+1}^{\infty} |I_k| \right) \leq 2\rho_1 \left(\lambda + \sum_{j=1}^{\infty} \frac{1}{\lambda^j} \right) |a| \\ &\leq 2\rho_1 (\lambda + 1) \text{per}(D). \end{aligned} \quad (12)$$

Putting (7) and (12) together yields

$$\begin{aligned} \text{per}(S_a) &= \text{per}(S_{\text{close}}) + \text{per}(S_{\text{far}}) \leq 2\rho_1 \left((\lambda + 1) \text{per}(D) + \frac{\text{esc}(S)}{\rho_2} \lambda \right) \\ &\leq 2\rho_1 \max \left(2, \frac{1}{\rho_2} \right) (\text{per}(D) + \text{esc}(S)) \lambda \leq \rho(C, D) (\text{per}(D) + \text{esc}(S)) \frac{\log n}{\log \log n}, \end{aligned}$$

for a suitable $\rho(C, D)$ depending on C and D , as required; here we set $\rho(C, D) = 8\rho_1 \max(2, 1/\rho_2)$.

Lower bound for squares. We first confirm the given lower bound for squares, i.e., we construct a packing S of $O(n)$ axis-aligned squares in the unit square $U = [0, 1]^2$ with total perimeter $\Omega((\text{per}(U) + \text{esc}(S)) \log n / \log \log n)$.

Let $n \geq 4$, and put $\lambda = \lfloor \log n / \log \log n \rfloor / 2$. We arrange each square $C_i \in S$ such that $\text{per}(C_i) = \lambda \text{esc}(C_i)$. We construct S as the union of λ subsets $S = \bigcup_{j=1}^{\lambda} S_j$, where S_j is a set of congruent squares, at the same distance from the bottom side of U .

Let S_1 be a singleton set consisting of one square of side length $1/4$ (and perimeter 1) at distance $1/\lambda$ from the bottom side of U . Let S_2 be a set of 2λ squares of side length $1/(4 \cdot 2\lambda)$ (and perimeter $1/(2\lambda)$), each at distance $1/(2\lambda^2)$ from the bottom side of U . Note that these squares lie strictly below the first square in S_1 , since $1/(8\lambda) + 1/(2\lambda^2) < 1/\lambda$. The total length of the vertical projections of the squares in S_2 is $2\lambda \cdot 1/(8\lambda) = 1/4$.

Similarly, for $j = 3 \dots, \lambda$, let S_j be a set of $(2\lambda)^{j-1}$ squares of side length $\frac{1}{4 \cdot (2\lambda)^{j-1}}$ (and perimeter $1/(2\lambda)^{j-1}$), each at distance $1/(2^{j-1}\lambda^j)$ from the bottom side of U . These squares lie strictly below any square in S_{j-1} ; and the total length of their vertical projections onto the x -axis is $(2\lambda)^{j-1} \cdot \frac{1}{4 \cdot (2\lambda)^{j-1}} = 1/4$.

The number of squares in $S = \bigcup_{j=1}^{\lambda} S_j$ is

$$\sum_{j=1}^{\lambda} (2\lambda)^{j-1} = \Theta\left((2\lambda)^{\lambda}\right) = O(n).$$

The total distance from the squares to the boundary of U is

$$\text{esc}(S) = \sum_{j=1}^{\lambda} (2\lambda)^{j-1} \frac{1}{2^{j-1}\lambda^j} = \lambda \frac{1}{\lambda} = 1.$$

The total perimeter of all squares in S is

$$4 \cdot \sum_{j=1}^{\lambda} \frac{1}{4} = \lambda = \Omega\left(\frac{\log n}{\log \log n}\right) = \Omega\left(\left(\text{per}(U) + \text{esc}(S)\right) \frac{\log n}{\log \log n}\right),$$

as required.

General lower bound. We now establish the lower bound in the general setting. Given a convex body C and a convex polygon D parallel to C , we construct a packing S of $O(n)$ positive homothets of C in D with total perimeter $\Omega((\text{per}(D) + \text{esc}(S)) \log n / \log \log n)$.

Let a be an arbitrary side of D . Assume w.l.o.g. that a is horizontal. Let U_C be the minimum axis-aligned square containing C . Clearly, we have $\frac{1}{2} \text{per}(U_C) \leq \text{per}(C) \leq \text{per}(U_C)$. We first construct a packing S_U of $O(n)$ axis-aligned squares in D such that for each square $U_i \in S_U$, $\text{esc}(U_i)$ equals the distance from the horizontal side a . We then obtain the packing S by inscribing a homothet C_i of C in each square $U_i \in S_U$ such that C_i touches the bottom side of U_i . Consequently, we have $\text{per}(S) \geq \text{per}(S_U)/2$ and $\text{esc}(S) = \text{esc}(S_U)$, since $\text{esc}(C_i) = \text{esc}(U_i)$ for each square $U_i \in S_U$.

It remains to construct the square packing S_U . Let $U(a)$ be a maximal axis-aligned square contained in D such that its bottom side is contained in a . S_U is a packing of squares in $U(a)$ that is homothetic with the packing of squares in the unit square U described previously. Put $\rho_1 = \text{per}(U(a))/\text{per}(U) = \text{per}(U(a))/4$. We have $\text{per}(S) \geq \frac{1}{4} \rho_1 \Omega\left(\left(\text{per}(U) + \text{esc}(S)\right) \frac{\log n}{\log \log n}\right)$, or

$$\text{per}(S) \geq \rho(C, D) \left((\text{per}(D) + \text{esc}(S)) \frac{\log n}{\log \log n} \right),$$

where $\rho(C, D)$ is a factor depending on C and D , as required. \square

Acknowledgment. We are grateful to Guangwu Xu for kindly showing us a short alternative derivation of the number theoretical sum in Equation (5). We also thank an anonymous reviewer for his very careful reading of the manuscript and his pertinent remarks.

References

- [1] T. M. Apostol, *Introduction to Analytic Number Theory*, Springer, New York, 1976.
- [2] S. Arora, Polynomial time approximation schemes for Euclidean traveling salesman and other geometric problems, *J. ACM* **45(5)** (1998), 753–782.
- [3] K. Bezdek, On a strong version of the Kepler conjecture, *Mathematika* **59(1)** (2013), 23–30.
- [4] M. Bern and D. Eppstein, Approximation algorithms for geometric problems, in *Approximation Algorithms for NP-hard Problems*, (D. S. Hochbaum, ed.), PWS Publishing Company, Boston, MA, 1997, pp. 296–345.
- [5] P. Brass, W. O. J. Moser, and J. Pach, *Research Problems in Discrete Geometry*, Springer, New York, 2005.
- [6] M. de Berg, J. Gudmundsson, M. J. Katz, C. Levcopoulos, M. H. Overmars, and A. F. van der Stappen, TSP with neighborhoods of varying size, *J. Algorithms* **57(1)** (2005), 22–36.
- [7] A. Dumitrescu and J. S. B. Mitchell, Approximation algorithms for TSP with neighborhoods in the plane, *J. Algorithms* **48(1)** (2003), 135–159.
- [8] A. Dumitrescu and C. D. Tóth, Minimum weight convex Steiner partitions, *Algorithmica* **60(3)** (2011), 627–652.
- [9] A. Dumitrescu and C. D. Tóth, The traveling salesman problem for lines, balls and planes, in *Proc. 24th ACM-SIAM Symposium on Discrete Algorithms*, 2013, SIAM, pp. 828–843.
- [10] K. R. Ford, Fractions, *The American Mathematical Monthly* **45(9)** (1938), 586–601.
- [11] A. Glazyrin and F. Morić, Upper bounds for the perimeter of plane convex bodies, *Acta Mathematica Hungarica* **142(2)** (2014), 366–383.
- [12] R. L. Graham, J. C. Lagarias, C. L. Mallows, A. R. Wilks, and C. H. Yan, Apollonian circle packings: geometry and group theory I: The Apollonian group, *Discrete Comput. Geom.* **34** (2005), 547–585.
- [13] T. C. Hales, The strong dodecahedral conjecture and Fejes Tóth’s conjecture on sphere packings with kissing number twelve, in *Discrete Geometry and Optimization*, vol. 69 of Fields Communications, 2013, Springer, 2013, pp. 121–132.
- [14] C. Levcopoulos and A. Lingas, Bounds on the length of convex partitions of polygons, in *Proc. 4th Annual Conference on Foundations of Software Technology and Theoretical Computer Science*, LNCS 181, 1984, Springer, pp. 279–295.
- [15] C. Mata and J. S. B. Mitchell, Approximation algorithms for geometric tour and network design problems, in *Proc. 11th ACM Symposium on Computational Geometry*, 1995, ACM, pp. 360–369.
- [16] J. S. B. Mitchell, A constant-factor approximation algorithm for TSP with pairwise-disjoint connected neighborhoods in the plane, in *Proc. 26th ACM Symposium on Computational Geometry*, 2010, ACM, pp. 183–191.